



10451 ROSELLE STREET, #300, SAN DIEGO, CA 92121  
TELEPHONE (858) 642-1988 • FAX (858) 642-1989  
WWW.ATSBIO.COM • ATS@ATSBIO.COM

**Antibody to Human c-Myc**  
**MOUSE MONOCLONAL**

**Catalog Number:** AB-407  
**Quantity:** 500 micrograms, 1 milligram  
**Format:** Lyophilized  
**Host:** synthetic peptides  
**Isotype:** IgG1  
**Clone:** NYRhc-myc

**Background:**

c-Myc is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of c-Myc have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of this gene.

**Specificity and Preparation:**

Mouse Anti-Human c-Myc is purified by Protein A. Protein concentration is 1 mg/ml in PBS (after reconstitution). This antibody was produced in BALB/c mice.

**Usage and Storage:**

Applications include Western blot (Titer: 1:2,000 will yield a strong band).

Reconstitute with sterile H<sub>2</sub>O. Mix gently, wash the sides of the vial and wait 30-60 seconds before use. In lyophilized form, for long periods, store at 4°C in a dry environment. After reconstitution, if not intended for use within a month, aliquot and store at -20°C. Antibody is stable for two years lyophilized, one month in solution at 4°C. Antibody is shipped lyophilized at ambient temperature. This material is provided for LABORATORY RESEARCH USE ONLY.