



10451 ROSELLE STREET, #300, SAN DIEGO, CA 92121
TELEPHONE (858) 642-1988 • FAX (858) 642-1989
WWW.ATSBIO.COM • ATS@ATSBIO.COM

**Antibody to Angiotensin II receptor (AT-1Br), affinity-purified
RABBIT POLYCLONAL**

Catalog Number: AB-N26AP
Quantity: 50 micrograms
Format: PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), with 0.1% sodium azide as preservative.
Host: Rabbit
Immunogen: peptide SSSAKKSASFFEVE (346-359) conjugated to keyhole limpet hemocyanin (KLH)

Background:

The Angiotensin II type 1 receptor (AT-1r) is the primary effector of Angiotensin II, a key regulator of blood pressure and fluid homeostasis. It is involved in pathogenesis of several cardiovascular diseases such as hypertension, cardiac hypertrophy and congestive heart failure. Angiotensin II interacts with two types of G-protein coupled membrane receptors, AT-1r (type 1) and AT-2r (type 2). AT-1 has three isoforms in rat: AT-1A (359 aa), AT-1B (359 aa), and AT-1C (177 aa). Rat AT-1r's are predicted to contain seven transmembrane domains. The N-terminus is predicted to be extracellular, while the C-terminus is predicted to be cytoplasmic. AT-1r's are expressed in the liver, kidney, aorta, lung, uterus, ovary, spleen, heart, adrenal and vascular smooth muscle.

Specificity and Preparation:

This antibody recognizes the AT-1B isoform of the Angiotensin II type 1 receptor in rat, and does not recognize isoform AT-1A or AT-2. The antisera was generated in rabbits by immunization with the peptide SSSAKKSASFFEVE (346-359) conjugated to keyhole limpet hemocyanin (KLH). Antisera was then affinity-purified by sequential passage through two affinity columns cross-linked with the carboxy terminal sequences of the rat AT-1B and AT-1A receptors. The AT-1B sequence was the same as that used for antibody production and the AT-1A peptide corresponded to amino acids 341-355. The antisera was retained by the first, but not the second column indicating recognition of the AT-1B receptor.

Usage and Storage:

Applications include immunolabeling (1:500).^{1,2} Store the antibody at -20°C for one year. Avoid repeated freezing and thawing. Gently spin down material before use; 5-10 seconds in a microfuge should be adequate.

References:

1. Huang J, Hara Y, Anrather J, Speth RC, Iadecola C, Pickel VM (2003) Angiotensin II subtype 1A (AT1A) receptors in the rat sensory vagal complex: Subcellular localization and association with endogenous angiotensin. *Neuroscience* 122(1):21-36.
2. Speth RC, Grove KL, Brownfield MS (2001) Immunohistochemical localization of AT-1A and AT-1B angiotensin II receptor subtypes in the rat adrenal. *Endocrine Soc Mtg*, Denver CO, Abstract.

To view protocol(s) for this and other products please visit: www.ATSBio.com/protocols