



10451 ROSELLE STREET, #300, SAN DIEGO, CA 92121  
TELEPHONE (858) 642-1988 • FAX (858) 642-1989  
WWW.ATSBIO.COM • ATS@ATSBIO.COM

**Antibody to NK-1 Receptor, Affinity-Purified**  
RABBIT POLYCLONAL

**Catalog Number:** AB-N33AP  
**Quantity:** 50 micrograms  
**Format:** PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), no preservative.  
**Host:** Rabbit  
**Immunogen:** Synthetic peptide corresponding to an amino acid sequence at the C-terminus of dog NK-1 receptor conjugated to bovine thyroglobulin with glutaraldehyde

**Background:**

The Neurokinin-1 (NK-1) receptor is a G-protein-coupled receptor characterized by seven transmembrane helices which preferentially binds the neuropeptide substance P. The NK-1 receptor (also known as the substance P receptor) plays a key role in pain and inflammation. Studies also implicate NK-1 receptors in depression and the growth of brain tumors.


**Specificity and Preparation:**

This antibody recognizes the NK-1 receptor in rat, dog, and human. This antibody was developed in rabbit using a synthetic peptide corresponding to an amino acid sequence at the C-terminus of dog NK-1 receptor conjugated to bovine thyroglobulin with glutaraldehyde. The peptide sequence has a high degree of homology to other species such as human, mouse, rat and guinea pig. The antibody is routinely tested by immunohistochemistry and flow cytometry.


**Usage and Storage:**

Applications include immunohistochemistry (ATS in-house, peroxidase substrate 1-3  $\mu\text{g/ml}$  and fluorescent 1-3  $\mu\text{g/ml}$ ), flow cytometry (ATS in-house, 3-10  $\mu\text{g/ml}$ ), immunoblotting (ATS in-house, western blot analysis 1-3  $\mu\text{g/ml}$ ), and ELISA (ATS in-house 1:1,000,000). Results may vary depending on protocol, tissue type etc; therefore the working dilutions should be determined by end user. Store the antibody at  $-20^{\circ}\text{C}$  for one year. Avoid repeated freezing and thawing. Gently spin down material before use; 5-10 seconds in a microfuge should be adequate.

**Antibody to NK-1 Receptor, Affinity-Purified  
RABBIT POLYCLONAL**



U-937 cells, human histiocytic lymphoma cell line, were used for a flow cytometry analysis with the NK-1r affinity-purified receptor antibody (lot #). Cells were treated with 4  $\mu\text{g}$  of the anti-NK-1r antibody and subsequently with anti-rabbit IgG-FITC. A 57% shift is seen as compared to the non-treated control.



Lane 1: Molecular weight standards (Invitrogen See-Blue).  
Lane 2: U-937 whole cell extract probed with anti-NK-1r affinity purified at 4  $\mu\text{g}/\text{ml}$ .  
Lane 3: Molecular weight standards (Invitrogen See-Blue).  
Lane 4: U-937 whole cell extract probed with anti-NK-1r affinity purified at 4  $\mu\text{g}/\text{ml}$  pre-absorbed with 25-fold molar excess of its cognate peptide

**To view protocol(s) for this and other products please visit: [www.ATSBio.com/protocols](http://www.ATSBio.com/protocols)**