



10451 ROSELLE STREET, #300, SAN DIEGO, CA 92121
TELEPHONE (858) 642-1988 • FAX (858) 642-1989
WWW.ATSBIO.COM • ATS@ATSBIO.COM

**Anti-Conjugated 5-Methoxytryptamine
RABBIT POLYCLONAL**

Catalog Number: AB-T078
Quantity: 50 microliters
Format: Lyophilized and reconstituted with deionized water / 50% glycerol.
Host: Rabbit
Isotype: IgG
Immunogen: synthetic 5-Methoxytryptamine conjugated to bovine serum albumin

Specificity and Preparation:

Antiserum previously preabsorbed on protein carriers and purified by ammonium sulfate precipitation.

This antibody targets conjugated 5-Methoxytryptamine.

Using a conjugate 5-Methoxytryptamine-Glutaraldehyde-BSA, antibody specificity was performed with an ELISA test by competition experiments with the following compounds:

COMPOUND	CROSS REACTIVITY §
5-Methoxytryptamine-G-BSA	1
5-Methoxytryptamine	1/20
5-Hydroxytryptamine-G-(Pc)	1/30
Tryptamine-G-(Pc)	1/30
5-Methoxytryptophan-G-(Pc)	1/40
Tryptophan	1/500
5-Hydroxytryptophan-G-(Pc)	1/1,900

G = Glutaraldehyde, BSA = bovine serum albumin

§ 5-Methoxytryptamine-G-BSA concentration/unconjugated or conjugated indolealkylamine concentration at half displacement.

Usage and Storage:

Applications include ELISA (1/1,000-1/5,000), immunocytochemistry, immunohistochemistry (1/1,000-1/5,000), immunoblotting (western blot 1/1,000-1/2,000). Fixation of tissue for use with these antibodies should be done with glutaraldehyde. The use of paraformaldehyde in conjunction with glutaraldehyde may improve staining results. Store the antibody at 4°C for one month or -20°C in undiluted aliquots for up to one year. Avoid repeated freezing and thawing. Gently spin down material before use; 5-10 seconds in a microfuge should be adequate.

Available Control(s): Corresponding antigen.

References:

1. Geffard M, Dulluc J, Heinrich-Rock AM (1985) Antisera against the indolealkylamines: tryptophan, 5-hydroxytryptophan, 5-hydroxytryptamine, 5-methoxytryptophan and 5-methoxytryptamine tested by an enzyme-linked immunosorbent assay method. *J Neurochem* 44:1221-1228.
2. Patel S, Dulluc J, Geffard M (1986) Comparison of serotonin and 5-methoxytryptamine immunoreactivity in rat raphe nuclei. *Histochem* 85:259-263.

To view protocol(s) for this product please visit: www.ATSBio.com/catalog/protocols