



10451 ROSELLE STREET, #300, SAN DIEGO, CA 92121
TELEPHONE (858) 642-1988 • FAX (858) 642-1989
WWW.ATSBIO.COM • ATS@ATSBIO.COM

**Anti-Conjugated 5,6-Dihydroxytryptamine
RABBIT POLYCLONAL**

Catalog Number: AB-T105
Quantity: 50 microliters
Format: Lyophilized and reconstituted with deionized water / 50% glycerol
Host: Rabbit
Isotype: IgG
Immunogen: Synthetic 5,6-Dihydroxytryptamine conjugated to bovine serum albumin

Specificity and Preparation:

Antiserum previously preabsorbed on protein carriers and purified by ammonium sulfate precipitation.

This antibody targets conjugated 5,6-Dihydroxytryptamine. **This antibody does not recognize free 5,6-dihydroxytryptamine.**

Using a conjugate 5,6-Dihydroxytryptamine-Glutaraldehyde-BSA, antibody specificity was performed with an ELISA test by competition experiments with the following compounds:

COMPOUND	CROSS REACTIVITY §
5,6-Dihydroxytryptamine-G-BSA	1
5,7-Dihydroxytryptamine-G-BSA	1/2,500
5-Hydroxytryptamine-G-BSA	1/>2,500
5-Hydroxytryptophan-G-BSA	1/>2,500

G = Glutaraldehyde, BSA = Bovine Serum Albumin

§ 5,6-Dihydroxytryptamine-G-BSA concentration/unconjugated or conjugated indolealkylamine concentration at half displacement.

Usage and Storage:

Applications include ELISA (1/1,000-1/5,000) and immunoblotting (western blot 1/1,000-1/2,000). Fixation of tissue for use with these antibodies should be done with glutaraldehyde. The use of paraformaldehyde in conjunction with glutaraldehyde may improve staining results. Store the antibody at 4°C for one month or -20°C in undiluted aliquots for up to one year. Avoid repeated freezing and thawing. Gently spin down material before use; 5-10 seconds in a microfuge should be adequate.

Available Control(s): 5,6-Dihydroxytryptamine-G-BSA

To view protocol(s) for this and other products please visit: www.ATSBio.com/protocols