



10451 ROSELLE STREET, #300, SAN DIEGO, CA 92121
TELEPHONE (858) 642-1988 • FAX (858) 642-1989
WWW.ATSBIO.COM • ATS@ATSBIO.COM

**Anti-Conjugated 5,7-Dihydroxytryptamine
RABBIT POLYCLONAL**

Catalog Number: AB-T106
Quantity: 50 microliters
Format: Lyophilized and reconstituted with deionized water / 50% glycerol.
Host: Rabbit
Isotype: IgG
Immunogen: synthetic 5,7-Dihydroxytryptamine conjugated to bovine serum albumin

Specificity and Preparation:

Antiserum previously preabsorbed on protein carriers and purified by ammonium sulfate precipitation.

This antibody targets conjugated 5,7-Dihydroxytryptamine.

Using a conjugate 5,7-Dihydroxytryptamine-Glutaraldehyde-BSA, antibody specificity was performed with an ELISA test by competition experiments with the following compounds:

| COMPOUND | CROSS REACTIVITY § |
|-------------------------------|---------------------------|
| 5,7-Dihydroxytryptamine-G-BSA | 1 |
| 5,6-Dihydroxytryptamine-G-BSA | 1/250 |
| 5-Hydroxytryptamine-G-BSA | 1/>50,000 |
| 5-Hydroxytryptophan-G-BSA | 1/>50,000 |

G = Glutaraldehyde, BSA = Bovine Serum Albumin

§ 5,7-Dihydroxytryptamine-G-BSA concentration/unconjugated or conjugated indolealkylamine concentration at half displacement.

Usage and Storage:

Applications include ELISA (1/1,000-1/5,000) and immunoblotting (western blot 1/1,000-1/2,000). Fixation of tissue for use with these antibodies should be done with glutaraldehyde. The use of paraformaldehyde in conjunction with glutaraldehyde may improve staining results. Store the antibody at 4°C for one month or -20°C in undiluted aliquots for up to one year. Avoid repeated freezing and thawing. Gently spin down material before use; 5-10 seconds in a microfuge should be adequate.

Available Control(s): Corresponding antigen.

To view example protocol(s) for this product please visit: www.ATSBio.com/catalog/protocols