

§

Anti-Conjugated NO-L-DOPA RAT POLYCLONAL

| Catalog Number: | AB-T121 |
|-----------------|---|
| Quantity: | 50 microliters |
| Format: | Lyophilized and reconstituted with deionized water / 50% glycerol |
| Host: | Rat |
| Immunogen: | Synthetic NO-L-DOPA conjugated to bovine serum albumin |

Specificity and Preparation:

Antiserum previously preabsorbed on protein carriers and purified by ammonium sulfate precipitation. This antibody targets conjugated NO-L-DOPA. **This antibody does not recognize free NO-L-DOPA.** Using a conjugate NO-L-DOPA-Glutaraldehyde-BSA, antibody specificity was performed with an ELISA test by competition experiments with the following compounds:

| COMPOUND | CROSS REACTIVITY |
|---|-------------------------|
| NO-L-DOPA-G-BSA | 1 |
| NO-Tyrosine-G-BSA | 1/40 |
| NO-Dopamine-G-BSA | 1/>50,000 |
| NO-Noradrenaline-G-BSA | 1/>50,000 |
| L-DOPA-G-BSA | 1/>50,000 |
| Cluteraldebude BSA - Devine Serum Albumin | |

G = Glutaraldehyde, BSA = Bovine Serum Albumin

§ NO-L-DOPA-G-BSA concentration/unconjugated or conjugated close-related compounds concentration at half displacement.

Usage and Storage:

Applications include ELISA (1/1,000-1/5,000), immunohistochemistry / immunocytochemistry (1/1,000 - 1/5,000), and immunoblotting (western blot 1/1,000-1/2,000). Fixation of tissue for use with these antibodies should be done with glutaraldehyde. The use of paraformaldehyde in conjunction with glutaraldehyde may improve staining results. Store the antibody at 4°C for one month or -20°C in undiluted aliquots for up to one year. Avoid repeated freezing and thawing. Gently spin down material before use; 5-10 seconds in a microfuge should be adequate.

Available Control(s): NO-L-DOPA-G-BSA

To view protocol(s) for this and other products please visit: www.ATSbio.com/protocols