

# Anti-Conjugated Glutathione RAT POLYCLONAL

Catalog Number: AB-T139
Quantity: 50 microliters

Format: Lyophilized and reconstituted with deionized water / 50% glycerol

**Host:** Rat **Isotype:** IgG

**Immunogen:** Synthetic glutathione conjugated to bovine serum albumin

# **Specificity and Preparation:**

Antiserum previously preabsorbed on protein carriers and purified by ammonium sulfate precipitation.

This antibody targets conjugated glutathione. This antibody does not recognize free glutathione.

Using a conjugate Glutathione-Glutaraldehyde-BSA, antibody specificity was performed with an ELISA test by competition experiments with the following compounds:

COMPOUND	CROSS REACTIVITY §
Glutathione-G-BSA	1
Cysteine-G-BSA	1/>10,000
Glutamate-G-BSA	1/>10,000
Glycine-G-BSA	1/>10,000

G = Glutaraldehyde, BSA = Bovine Serum Albumin

### **Usage and Storage:**

Applications include ELISA (1/2,000-1/5,000) and immunohistochemistry / immunocytochemistry (1/500 -1/1,000). Fixation of tissue for use with these antibodies should be done with glutaraldehyde. The use of paraformaldehyde in conjunction with glutaraldehyde may improve staining results. Store the antibody at 4°C for one month or -20°C in undiluted aliquots for up to one year. Avoid repeated freezing and thawing. Gently spin down material before use; 5-10 seconds in a microfuge should be adequate.

# Available Control(s): Glutathione-G-BSA

#### **References:**

1. Amara A, Coussemacq M, Geffard M. (1994) Antibodies to reduced glutathione. Brain Res 659(1-2):237 -42.

To view protocol(s) for this and other products please visit: www.ATSbio.com/protocols

<sup>§</sup> Glutathione-G-BSA concentration/unconjugated or conjugated analogs concentration at half displacement.