



NPY-SAP
TARGETED SAP CONJUGATE

*a tool for eliminating cells that express the Neuropeptide Y (NPY);
targeted via NPY, eliminated via saporin*

Catalog Number: IT-28
Quantity: 25 micrograms, 100 micrograms, 250 micrograms, 1 milligram
Format: PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), no preservative. Sterile-filtered.

Background: Targeted SAP conjugates are powerful and specific lesioning agents used in the technique known as Molecular Surgery. The ribosome-inactivating protein, saporin (from the seeds of the plant, *Saponaria officinalis*) is bound to a targeting agent (anything that is recognized on the cell surface and internalized). The targeted conjugate is administered to cells (*in vitro* or *in vivo*). The targeting agent seeks out and binds to its target on the cell surface. The conjugate is internalized, saporin breaks away from the targeting agent, and inactivates the ribosomes which causes protein inhibition and, ultimately, cell death. Cells that do not have the cell surface marker are not affected.

Neuropeptide Y (NPY) receptor is the most abundant neuropeptide in the brain and is involved in many processes from prenatal to mature animals. It promotes the proliferation of postnatal neuronal precursor cells and exhibits a diverse range of important physiologic activities, including effects on psychomotor activity, food intake, regulation of central endocrine secretion, and potent vasoactive effects on the cardiovascular system. Neuropeptide Y family of peptides signal through a family of G protein-coupled receptors present in the brain and sympathetic neurons. These NPY receptors recognize NPY and related peptides. NPY-SAP specifically eliminates cells expressing the NPY receptor. It is not suitable for retrograde transport.

Specificity & Preparation: This immunotoxin (molecular weight 34 kDa) recognizes cells expressing the Neuropeptide Y (NPY) receptor in rat and mouse. NPY-SAP is a chemical conjugate of neuropeptide Y and the ribosome-inactivating protein, saporin.

Usage: NPY-SAP specifically eliminates cells expressing the NPY receptor. Not suitable for retrograde transport. **There may be lot-to-lot variation in material; working dilutions must be determined by end user. If this is a new lot, you must assess the proper working dilution before beginning a full experimental protocol.**

Storage: Gently spin down material 5-10 seconds in a microfuge before use. Store the material in undiluted aliquots at -20°C for 1-2 months. For longer term storage store the material at -80°C . Material should be aliquoted to a convenient volume and quantity to avoid repeated freezing and thawing that can damage the protein content. Under these conditions, the material has a very stable shelf-life. Thawing should be done at room temperature or on ice. The thawed solution should remain on ice until use.

Do not use a reducing agent (such as dithiothreitol, beta-mercaptoethanol or ascorbic acid) with this material. It will inactivate the toxin.

This material is an extremely potent cytotoxin. Handling should be done by experienced personnel. Gloves and safety glasses are required when handling this product. Care in disposal is mandatory; autoclaving or exposure to 0.2 M sodium hydroxide will inactivate the material. All labware that comes into contact with this material should be likewise treated.



NPY-SAP
TARGETED SAP CONJUGATE

**Selected References:**

1. Li AJ, Ritter S (2003) 2-Deoxy-D-Glucose (2DG) increases NPY mRNA expression in hindbrain neurons. *Soc Neurosci Mtg, New Orleans, LA*, Abstract #615.7.
2. Bugarith K, Ritter S, Dinh T (2001) Hyperphagia and obesity results from the injection of the immunotoxin neuropeptide Y (NPY)-saporin (NPY-SAP) into the paraventricular hypothalamus (PVH) of rats. *Soc Neurosci Mtg, San Diego, CA*, Abstract #948.2.

Scan to view
all product
references.

Control(s): Blank-SAP

Safety:

Good laboratory technique must be employed for safe handling of this product. This requires observation of the following practices:

1. Wear appropriate laboratory attire, including lab coat, gloves and safety glasses.
2. Do not pipet by mouth, inhale, ingest or allow product to come into contact with open wounds. Wash thoroughly any part of the body which comes into contact with the product.
3. Avoid accidental autoinjection by exercising extreme care when handling in conjunction with any injection device.
4. This product is intended for research use by qualified personnel only. It is not intended for use in humans or as a diagnostic agent. Advanced Targeting Systems is not liable for any damages resulting from the misuse or handling of this product.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols