



10451 ROSELLE STREET, #300, SAN DIEGO, CA 92121
TELEPHONE (858) 642-1988 • FAX (858) 642-1989
WWW.ATSBIO.COM • ATS@ATSBIO.COM

**Anti-Anti-Conjugated L-Glutamate
RABBIT POLYCLONAL
AB-T021**

Example of Immunochemistry protocol:

Perfusion protocol for Adult male Sprague Dawley (weight around 0.5 kg):

1. The animals can be deeply anaesthetized (for example with urethane-0.5-1.5g/kg, intraperitoneal).
2. Perfused via the ascending aorta with 50 ml of NaCl 9g/l (Heparinized) and pass through the system 800-1000 ml of cold 4% paraformaldehyde (Merck) in 0.1 M PB, pH 7.2-7.4, (ten minutes).
3. Dissect out the organs and place in a solution of 4% paraformaldehyde in 0.1M PB, pH 7.2, at 4°C for twelve to sixteen hours.

Immunohistochemistry

1. In order to avoid possible interference with endogenous peroxidase, free-floating sections will be treated with distilled water containing NH₃ (20%), H₂O₂ (30%) and NaOH (1%) for 20 min (other method is using a solution with 33% of H₂O₂ and 66% of methanol).
2. Then, wash the sections for 20 min in 0.15 M phosphate-buffered saline (PBS) (pH 7.2)
3. Pre-incubate for 30 min in PBS containing 2-10% (variable to adjust) of normal horse serum and 0.3% of Triton X-100 (mixed solution).
4. Incubate at room temperature (1h 30min) and overnight at 4°C in the same mixed solution containing the diluted anti-anti conjugated L-Glutamate antibodies (1/1,000-1/5,000).
5. Then, the sections will be wash in PBS (30 min).
6. After that we will incubate for 60 min at room temperature with biotinylated anti-(species) immunoglobulin (Vector) diluted 1/200 in PBS.
7. Wash during 30 min with PBS.
8. Sections will be incubated for 1 h with a 1/100 diluted avidin-biotin-peroxidase complex (Vectastain) in the mixed solution.
9. After that we will wash the sections in PBS (30 min)
10. Wash with Tris-HCl buffer (pH 7.6)(10 min).
11. The tissue-bound peroxidase will be developed with H₂O₂ using 3, 3' diaminobenzidine as chromogen.
12. Finally the sections will be rinsed with PBS and coverslipped with PBS/Glycerol (1/1).