

**Alexa488-labeled Antibody to Saporin, Affinity-Purified**
GOAT POLYCLONAL

Catalog Number: AB-15AP-FLA
Quantity: 50 micrograms
Format: 50% PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), 50% glycerol; no preservative.
Host: Goat
Isotype: IgG
Immunogen: Saporin

Background: Saporin is obtained from the seeds of the Soapwort plant (*Saponaria officinalis*), a plant that grows wildly in Britain and other parts of Europe. Saporin is a plant enzyme with N-glycosidase activity that deurinates a specific nucleotide in the ribosomal RNA 28S, thus irreversibly blocking protein synthesis. It belongs to the well-characterized family of ribosome-inactivating proteins (RIPs). There are two types of RIPs: type I, which are much less cytotoxic due to the lack of the B chain and type II, which are distinguished from type I RIPs by the presence of the B chain and their ability to enter cells on their own. However, type I RIPs can still be internalized by fluid-phase endocytosis. Upon internalization, the ribosomes are inactivated, resulting in cell death.

Specificity & Preparation: This antibody recognizes saporin. Saporin was used as the immunogen. The antibody was affinity-purified against saporin attached to a CnBr-Sepharose support column. It has been conjugated to the fluorescent dye Alexa488. The antibody is routinely tested by Western blot.

Usage: Applications include immunoblotting (ATS in-house; 1:1,000).

Storage: Gently spin down material 5-10 seconds in a microfuge before use. The material can be handled safely using normal laboratory precautions. Store the antibody at -20°C for up to one year.



Scan to view
all product
references.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols