

**Alexa488-labeled Antibody to Human CD25 (Tac, IL-2 receptor)**  
**MOUSE MONOCLONAL**

**Catalog Number:** AB-18-FLA  
**Quantity:** 50 micrograms  
**Format:** 50% PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), 50% glycerol; no preservative.  
**Host:** Mouse  
**Isotype:** IgG<sub>2A</sub>  
**Immunogen:** Cultured human T cells

**Background:** The monoclonal CD25 antibody inhibits the proliferation of Interleukin-2 (IL-2)-dependent human continuous T-cell lines. It blocks the human T-cell receptor for IL-2 by binding to the alpha-chain of the IL-2 receptor. In antigen-induced T-cell proliferative assays, the addition of highly purified IL-2 reverses the inhibitory effects of anti-Tac.

**Specificity & Preparation:** This antibody recognizes the human Interleukin-2 (IL-2) receptor. Cultured human T-cells were used as immunogen. This antibody was produced in tissue culture supernatants. It has been conjugated to the fluorescent dye Alexa488. The antibody is routinely tested by flow cytometry.

**Usage:** Applications include cytotoxic killer cell activity (50  $\mu$ l culture supernatant),<sup>1</sup> proliferation inhibition,<sup>2</sup> and flow cytometry (1:10,000).<sup>3</sup>

**Storage:** Gently spin down material 5-10 seconds in a microfuge before use. The material can be handled safely using normal laboratory precautions. Store the antibody at -20°C for up to one year.



Scan to view  
all product  
references.

**Selected References:**

1. Uchiyama T, Broder S, Waldmann TA (1981) A monoclonal antibody (anti-Tac) reactive with activated and functionally mature human T cells. I. Production of anti-Tac monoclonal antibody and distribution of Tac (+) cells. *J Immunol* 126(4):1393-1397.
2. Uchiyama T, Nelson DL, Fleisher TA, Waldmann TA (1981) A monoclonal antibody (anti-Tac) reactive with activated and functionally mature human T cells. II. Expression of Tac antigen on activated cytotoxic killer T cells, suppressor cells, and on one of two types of helper T cells. *J Immunol* 126(4):1398-1403.
3. Depper JM, Leonard WJ, Robb RJ, Waldmann TA, Greene WC (1983) Blockade of the interleukin-2 receptor by anti-Tac antibody: Inhibition of human lymphocyte activation. *J Immunol* 131(2):690-696.

To view protocol(s) for this and other products please visit: [www.ATSBio.com/library/protocols](http://www.ATSBio.com/library/protocols)