

**Antibody to Mouse CD25 (Tac, IL-2 receptor)**
RAT MONOCLONAL

Catalog Number: AB-19
Quantity: 50 micrograms
Format: PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), no preservative.
Host: Rat
Isotype: IgG₁
Clone: PC61 5.3
Immunogen: Cultured mouse B6.1 cytotoxic T-cells

Background: Interleukin-2 receptors are located on the surface of T-cells and function in clonal expansion of the activated T-cell. The monoclonal CD-25 antibody inhibits the proliferation of Interleukin-2 (CD25)-dependent murine cell lines. It blocks the murine T-cell receptor for IL-2 by binding to the alpha-chain of IL-2 receptors. This antibody has also been shown to inhibit IL-2 binding to both low and high affinity IL-2 receptors.

Specificity & Preparation: This antibody recognizes the murine interleukin-2 (CD25) receptor. Cultured mouse B6.1 cytotoxic T-cells were used as immunogen. This antibody was produced in tissue culture supernatants.

Usage: Applications include proliferation inhibition (0.01-1.0 $\mu\text{g/ml}$),^{2,3} flow cytometry (2 μg),¹ immunoprecipitation,³ targeting (targeting agent in Anti-CD25-SAP, Cat. #IT-29).

Storage: Store the antibody at 4°C for one month or -20°C in undiluted aliquots for one year. Avoid repeated freezing and thawing. Gently spin down material 5-10 seconds in a microfuge before use.



Scan to view
all product
references.

Selected References:

1. Rusterholz C, Henrioud PC, Nabholz M (1999) Interleukin-2 (IL-2) regulates the accessibility of the IL-2-responsive enhancer in the IL-2 receptor alpha gene to transcription factors. *Mol Cell Biol* 19(4):2681-2689.
2. Lowenthal JW, Corthesy P, Tougne C, Lees R, MacDonald HR, Nabholz M (1985) High and low affinity IL 2 receptors: analysis by IL 2 dissociation rate and reactivity with monoclonal anti-receptor antibody PC61. *J Immunol* 135(6):3988-3994.
3. Lowenthal JW, Zubler RH, Nabholz M, MacDonald HR (1985) Similarities between interleukin-2 receptor number and affinity on activated B and T lymphocytes. *Nature* 315 (6021):669-672.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols