



### Antibody to Saporin, Affinity-Purified RABBIT POLYCLONAL

**Catalog Number:** AB-41AP  
**Quantity:** 50 micrograms  
**Format:** PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), no preservative.  
**Host:** Rabbit  
**Isotype:** IgG  
**Immunogen:** Saporin

**Background:** Saporin is obtained from the seeds of the Soapwort plant (*Saponaria officinalis*), a plant that grows wild in Britain and other parts of Europe. Saporin is a plant enzyme with N-glycosidase activity that dephosphorylates a specific nucleotide in the ribosomal RNA 28S, thus irreversibly blocking protein synthesis. It belongs to the well-characterized family of ribosome-inactivating proteins (RIPs). There are two types of RIPs: type I, which are much less cytotoxic due to the lack of the B chain and type II, which are distinguished from type I RIPs by the presence of the B chain and their ability to enter cells on their own. However, type I RIPs can still be internalized by fluid-phase endocytosis. Upon internalization, the ribosomes are inactivated, resulting in cell death.

**Specificity & Preparation:** This antibody recognizes saporin. The antibody was first protein-A-purified followed by affinity-purification against saporin attached to a CnBr-Sepharose support column. This antibody is routinely tested by Western Blot.

**Usage:** Applications include immunoblotting (Western Blot ATS in-house; 1:1,000).

**Storage:** Store the antibody at 4°C for one month or -20°C in undiluted aliquots for one year. Avoid repeated freezing and thawing. Gently spin down material 5-10 seconds in a microfuge before use.

To view protocol(s) for this and other products please visit: [www.ATSBio.com/library/protocols](http://www.ATSBio.com/library/protocols)

