



Antibody to Nerve Growth Factor (p75) Receptor, Affinity-Purified RABBIT POLYCLONAL

Catalog Number: AB-N01AP
Quantity: 50 micrograms
Format: PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), no preservative.
Host: Rabbit
Immunogen: extracellular fragment from the mouse p75 receptor (amino acids 43-161)

Background: The p75 neurotrophin receptor (p75^{NTR}), also known as the low affinity nerve growth factor receptor, binds nerve growth factor, brain-derived neurotrophic factor, neurotrophin-3 and neurotrophin-4 with varying specificities. The p75^{NTR} plays an important role in neurotrophic factor signaling and has been shown to modulate the susceptibility of selective cellular populations to programmed cell death.

Specificity & Preparation: This antibody recognizes p75^{NTR} in mouse. The antisera was developed in rabbit using an extracellular fragment from the mouse p75 receptor (amino acids 43-161). The antibody was affinity-purified using the extracellular domain of p75. The antibody is routinely tested by flow cytometry.

Usage: Applications include immunohistochemistry (paraffin sections; 1:100)¹ and flow cytometry (ATS in-house; 1:1,000),² targeting (targeting agent in mu p75-SAP, Cat. #IT-16).

Storage: Store the antibody at -20°C for one year. Avoid repeated freezing and thawing. Gently spin down material 5-10 seconds in a microfuge before use.





Selected References:

1. Rock JR, Onaitis MW, Rawlins EL, Lu Y, Clark CP, Xue Y, Randell SH, Hogan BL (2009) Basal cells as stem cells of the mouse trachea and human airway epithelium. *Proc Natl Acad Sci U S A* 106(31):12771-12775.
2. Lopez-Coviella I, Follettie MT, Mellott TJ, Kovacheva VP, Slack BE, Diesl V, Berse B, Thies RS, Blusztajn JK (2005) Bone morphogenetic protein 9 induces the transcriptome of basal forebrain cholinergic neurons. *Proc Natl Acad Sci U S A* 102(19):6984-6989.

Scan to view
all product
references.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols



Immunofluorescent staining of NG6 cells* with anti-mu p75. Cells were fixed with paraformaldehyde and blocked prior to staining. Primary was added at 10 µg/ml followed by goat anti-rabbit-FITC (A) and goat anti-rabbit-Cy3 (B) secondary at 15 µg/ml (5 µg/ml for DAPI nuclear staining).

Images were obtained using a 40x (A) and 60X (B) objective and a fluorescent microscope. NGFr staining is represented in green (A) and red (B) and nuclear staining is represented in blue.

*clone of the NG108-15 fusion of mouse neuroblastoma and rat glioblastoma⁺