

**Biotin-labeled Antibody to Mac-1 (CD11b)**
RAT MONOCLONAL (IGG_{2B})

Catalog Number: AB-N05-BT
Quantity: 50 micrograms
Format: PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), no preservative. Sterile-filtered.
Host: Rat
Isotype: IgG_{2b}
Clone: CD11b
Immunogen: B10 mouse spleen cells enriched for T-lymphocytes

Background: CD11b is an alpha subunit of Mac-1, also known as CR3. CD11b is the receptor for the C3bi fragment of complement. This receptor is involved in bacterial phagocytosis. A reduction in neutrophil CD11b expression after severe traumatic injury correlates with increased septic complications. CD11b is a component of integrins, important for adhesion of neutrophils to surfaces. Mac-1 exists as a chemoattractant activation-dependent molecule that undergoes a conformational change upon stimulation. Expression of new epitopes on Mac-1 can be detected after activation by specific reporter monoclonal antibodies. Until stimulation occurs, Mac-1 remains in a resting, non-adhesive state. Activation of Mac-1 may play a role during neutrophil recruitment to the inflamed site.

Specificity & Preparation: This antibody recognizes human and mouse Mac-1 (CD11b). The hybridoma was formed by the fusion of mouse myeloma NS1 cells with spleen cells from rats immunized with B10 mouse spleen cells enriched for T-lymphocytes. It has been conjugated to biotin via an amide bond. The antibody is routinely tested by flow cytometry.

Usage: Applications include immunoprecipitation,¹ flow cytometry (50 μ l),² cytotoxicity assay (EC₅₀=16 pM),³ and binding assay (5 and 50 μ l of culture supernatant).³

Storage: Store the antibody at -20°C for one year. Avoid repeated freezing and thawing. Gently spin down material 5-10 seconds in a microfuge before use.



Scan to view
all product
references.

Selected References:

1. Sanchez-Madrid F, Simon P, Thompson S, Springer TA. (1983) Mapping of antigenic and functional epitopes on the alpha- and beta-subunits of two related mouse glycoproteins involved in cell interactions, LFA-1 and Mac-1. *J Exp Med* 158:586-602.
2. Springer T, Galfre G, Secher DS, Milstein C. (1979) Mac-1: a macrophage differentiation antigen identified by monoclonal antibody. *Eur J Immunol* 9:301-306.
3. Springer, T., Galfre, G., Secher, D.S., Milstein, C. (1978) Monoclonal xenogeneic antibodies to murine cell surface antigens: identification of novel leukocyte differentiation antigens. *Eur J Immunol* 8:539-551.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols