



Biotin-labeled Antibody to Somatostatin Receptor-4 (SSTR4), Affinity-Purified RABBIT POLYCLONAL

| Catalog Number: | AB-N21AP-BT |
|-----------------|--|
| Quantity: | 50 micrograms |
| Format: | PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium |
| | Phosphate; 0.01 M Sodium Phosphate; pH 7.4), no preservative. |
| Host: | Rabbit |
| Immunogen: | peptide corresponding to the extracellular domain of rat SSTR4 conjugated to keyhole limpet hemocyanin (KLH) |

Background: Somatostatin Receptor-4 is one of five receptor subtypes termed SSTR1-5. They are G-proteincoupled receptors characterized by seven transmembrane domains with an extracellular amino terminal domain and an intracellular carboxy terminus. These receptors function in the regulation of numerous physiological processes such as the secretion of insulin, glucagon, and growth hormone, as well as cell growth induced by neuronal excitation in both the central and peripheral nervous systems. Somatostatin receptors are activated via somatostatin secretion in nerve and endocrine cells.

Specificity & Preparation: This antibody was raised against rat somatostatin receptor-4 (SSTR4) and recognizes SSTR4 in rat. The SSTR antisera was developed in rabbit using a peptide (Cat. #PR-07) corresponding to the extracellular domain conjugated to keyhole limpet hemocyanin (KLH) for immunization. Antisera was then affinity-purified with the peptide utilized for immunization. It has been conjugated to biotin via an amide bond. The antibody is routinely tested by immunoblotting.

Usage: Applications include immunoblotting (ATS in-house) using a dilution of 1:500-1:1,000 and ELISA (ATS in-house).

Storage: Store the antibody at -20°C for one year. Avoid repeated freezing and thawing. Gently spin down material 5-10 seconds in a microfuge before use.



Scan to view all product references.

Control(s): SSTr4 peptide

To view protocol(s) for this and other products please visit: www.ATSbio.com/library/protocols