

**Antibody to NK-1 Receptor, Affinity-Purified**  
RABBIT POLYCLONAL

**Catalog Number:** AB-N33AP  
**Quantity:** 50 micrograms  
**Format:** PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), no preservative.  
**Host:** Rabbit  
**Immunogen:** Synthetic peptide corresponding to an amino acid sequence at the C-terminus of dog NK-1 receptor conjugated to bovine thyroglobulin with glutaraldehyde

**Background:** The Neurokinin-1 (NK-1) receptor is a G-protein-coupled receptor characterized by seven transmembrane helices which preferentially binds the neuropeptide substance P. The NK-1 receptor (also known as the substance P receptor) plays a key role in pain and inflammation. Studies also implicate NK-1 receptors in depression and the growth of brain tumors.

**Specificity & Preparation:** This antibody recognizes the NK-1 receptor in rat, dog, and human. This antibody was developed in rabbit using a synthetic peptide corresponding to an amino acid sequence at the C-terminus of dog NK-1 receptor conjugated to bovine thyroglobulin with glutaraldehyde. The peptide sequence has a high degree of homology to other species such as human, mouse, rat and guinea pig. The antibody is routinely tested by immunohistochemistry and flow cytometry.

**Usage:** Applications include immunohistochemistry (ATS in-house, peroxidase substrate 1-3  $\mu\text{g/ml}$  and fluorescent 1-3  $\mu\text{g/ml}$ ), flow cytometry (ATS in-house, 3-10  $\mu\text{g/ml}$ ), immunoblotting (ATS in-house, western blot analysis 1-3  $\mu\text{g/ml}$ ), and ELISA (ATS in-house 1:1,000,000). Results may vary depending on protocol, tissue type etc; therefore the working dilutions should be determined by end user.

**Storage:** Store the antibody at  $-20^{\circ}\text{C}$  for one year. Avoid repeated freezing and thawing. Gently spin down material 5-10 seconds in a microfuge before use.




**Antibody to NK-1 Receptor, Affinity-Purified  
RABBIT POLYCLONAL**



Scan to view  
all product  
references.

To view protocol(s) for this and other products please visit: [www.ATSBio.com/library/protocols](http://www.ATSBio.com/library/protocols)



Immunoblot on U-937 (human histiocytic lymphoma) whole cell extract. Data shows extract probed with anti-NK1r versus extract probed with anti-NK1r pre-absorbed with molar excess of its cognate peptide.  
Lane 1: Molecular weight standards (Invitrogen See-Blue).  
Lane 2: U-937 whole cell extract probed with anti-NK-1r affinity purified at 4  $\mu$ g/ml.  
Lane 3: Molecular weight standards (Invitrogen See-Blue).  
Lane 4: U-937 whole cell extract probed with anti-NK-1r affinity purified at 4  $\mu$ g/ml pre-absorbed with 25-fold molar excess of its cognate peptide