



Antibody to Somatostatin Receptor-1 (SSTR1) MOUSE MONOCLONAL

Catalog Number: AB-N35
Quantity: 50 micrograms
Format: PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), no preservative.
Host: Mouse
Isotype: IgM Kappa
Clone: (15F10) 2D7
Immunogen: peptide corresponding to the extracellular domain of SSTR1 conjugated to keyhole limpet hemocyanin (KLH)

Background: Somatostatin Receptor-1 is one of the five subtypes termed SSTR1-5. They are G-protein-coupled receptors characterized by seven transmembrane helices with an extracellular amino terminal domain and an intracellular carboxy terminus. These receptors function in the regulation of numerous physiological processes such as the secretion of insulin, glucagon, and growth hormone, as well as cell growth induced by neuronal excitation in both the central and peripheral nervous system. Somatostatin receptors are activated via somatostatin secreted by nerve and endocrine cells.

Specificity & Preparation: This antibody was raised against rat somatostatin receptor-1 (SSTR1) and recognizes SSTR1 in human and rat. The SSTR1 monoclonal antibody was developed using a peptide corresponding to the extracellular domain of SSTR1 conjugated to keyhole limpet hemocyanin (KLH). This antibody is routinely tested by flow cytometry.

Usage: Applications include immunohistochemistry and immunocytochemistry (ATS in-house, 2-10 $\mu\text{g/ml}$), flow cytometry (ATS in-house, 2-10 $\mu\text{g/ml}$), immunoblotting (ATS in-house, western blot 2-10 $\mu\text{g/ml}$), and ELISA (ATS in-house, 1:125,000). The final working dilutions should be determined by end user.

Storage: Store the antibody at -20°C for one year. Gently spin down material 5-10 seconds in a microfuge before use.




Selected References:

1. Moller LN, Stidsen CE, Hartmann B, Holst JJ (2003) Somatostatin receptors. *Biochim Biophys Acta* 1616(1):1-84. Review.

Scan to view
all product
references.

Control(s): SSTR1 peptide

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols



FLOW CYTOMETRY ANALYSIS OF CHO CELLS EXPRESSING SSTR1 USING ANTI-SSTR1 MONOCLONAL ANTIBODY
CHO (Chinese Hamster Ovary) cells were transfected with rat SSTR1 (lot #49-48) and treated with either 2 $\mu\text{g/ml}$ of SSTR1 monoclonal antibody or 2 $\mu\text{g/ml}$ of normal mouse IgM (negative control) followed by anti-mouse IgM-FITC. A 74% shift was seen as compared to the negative control.