

**Cy3-labeled Antibody to Nerve Growth Factor (p75) Receptor (192 IgG)**  
MOUSE MONOCLONAL

**Catalog Number:** AB-N43-FL3  
**Quantity:** 50 micrograms  
**Format:** 50% PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), 50% glycerol; no preservative.  
**Host:** Mouse  
**Isotype:** IgG<sub>1</sub>  
**Clone:** 192

**Background:** 192-IgG is the antibody to the p75 neurotrophin receptor (p75<sup>NTR</sup>). The p75<sup>NTR</sup>, also known as the low affinity nerve growth factor receptor, binds nerve growth factor, brain-derived neurotrophic factor, neurotrophin-3 and neurotrophin-4 with varying specificities. The p75<sup>NTR</sup> plays an important role in neurotrophic factor signaling and has been shown to modulate the susceptibility of selective cellular populations to programmed cell death.

**Specificity & Preparation:** This fluorescent conjugate recognizes p75 receptor-positive cells in rat. It was prepared using mouse monoclonal antibody 192-IgG conjugated to Cy3. This product is routinely tested by flow cytometry.

**Usage:** Applications include flow cytometry (ATS in-house; 4  $\mu$ g/10<sup>6</sup> cells per 200  $\mu$ l), immunofluorescence,<sup>1</sup> immunocytochemistry (paraffin; 10-20  $\mu$ g/ml TBS).<sup>2</sup> This product can also be used as a transport agent.<sup>2</sup>

**Storage:** Store at -20°C for one year. Gently spin down material 5-10 seconds in a microfuge before use. The material can be handled safely using normal laboratory precautions.



**Cy3-labeled Antibody to Nerve Growth Factor (p75) Receptor (192 IgG)  
MOUSE MONOCLONAL**




Scan to view all product references.


**Selected References:**

1. Wu M, Shanabrough M, Leranath C, Alreja M (2000) Cholinergic excitation of septohippocampal GABA but not cholinergic neurons: implications for learning and memory. *J Neurosci* 20(10):3900-3908.
2. Hartig W, Seeger J, Naumann T, Brauer K, Bruckner G. (1998) Selective *in vivo* fluorescence labelling of cholinergic neurons containing p75(NTR) in the rat basal forebrain. *Brain Res* 808 (2):155-165.

To view protocol(s) for this and other products please visit: [www.ATSBio.com/library/protocols](http://www.ATSBio.com/library/protocols)



C6/9 cells, a rat glioma cell line, were labeled with 192-IgG conjugated to Cy3 dye at various concentrations, and allowed to incubate at 4°C for one hour. Unlabeled antibody was used as a control. Cells were then washed, and samples analyzed on a BD FACScan. Data was processed using CellQuest software. The conjugated 192 IgG-Cy3 demonstrated a 43% positive shift versus the control.



HEK-293 cells, transfected with the p75 neurotrophin receptor were labeled with 192-IgG conjugated to Cy3 at 20 mg/ml and incubated at 37°C for 24 hours. Cells were then washed with PBS and analyzed on a fluorescent microscope under 20X magnification using a Y3 Leica filter cube.