

**Alexa488-labeled Antibody to Nerve Growth Factor (p75) Receptor (192 IgG)**  
**MOUSE MONOCLONAL**

**Catalog Number:** AB-N43-FLA  
**Quantity:** 50 micrograms  
**Format:** 50% PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), 50% glycerol; no preservative.  
**Host:** Mouse  
**Isotype:** IgG<sub>1</sub>  
**Clone:** 192

**Background:** 192-IgG is the antibody to the p75 neurotrophin receptor (p75<sup>NTR</sup>). The p75<sup>NTR</sup>, also known as the low affinity nerve growth factor receptor, binds nerve growth factor, brain-derived neurotrophic factor, neurotrophin-3 and neurotrophin-4 with varying specificities. The p75<sup>NTR</sup> plays an important role in neurotrophic factor signaling and has been shown to modulate the susceptibility of selective cellular populations to programmed cell death.

**Specificity & Preparation:** This fluorescent conjugate recognizes p75 receptor-positive cells in rat. It was prepared using mouse monoclonal antibody 192-IgG conjugated to Alexa 488. This product is routinely tested by flow cytometry.

**Usage:** Applications include flow cytometry (ATS in-house; 1 µg/10<sup>6</sup> cells per 200 µl), fluorescence spectroscopy,<sup>1</sup> fluorescence anisotropy,<sup>1</sup> epifluorescence microscopy (100 nM),<sup>2</sup> qualitative probe of binding to membrane-bound IgG's and the cytoskeleton (0.1 µM).<sup>3</sup>

**Storage:** Store at -20°C for one year. Gently spin down material 5-10 seconds in a microfuge before use. The material can be handled safely using normal laboratory precautions.



**Alexa488-labeled Antibody to Nerve Growth Factor (p75) Receptor (192 IgG)**  
**MOUSE MONOCLONAL**



Scan to view  
all product  
references.

**Selected References:**

1. Harikumar KG, Pinon DI, Wessels WS, Prendergast FG, Miller LJ (2002) Environment and mobility of a series of fluorescent reporters at the amino terminus of structurally related peptide agonists and antagonists bound to the cholecystokinin receptor. *J Biol Chem* 277(21):18552-18560.
2. Bennett VJ, Simmons MA (2001) Analysis of fluorescently labeled substance P analogs: binding, imaging and receptor activation. *BMC Chem Biol* 1(1):1.
3. Salhany JM, Cordes KS, Sloan RL (2001) Characterization of immunoglobulin binding to isolated human erythrocyte membranes: evidence for selective, temperature-induced binding of naturally occurring autoantibodies to the cytoskeleton. *Biochim Biophys Acta* 1511(1):168-180.

To view protocol(s) for this and other products please visit: [www.ATSBio.com/library/protocols](http://www.ATSBio.com/library/protocols)