



6352 CORTE DEL ABETO, STE B
CARLSBAD, CA 92011 USA
01.858.642.1988 • WWW.ATSBIO.COM

Anti-Conjugated Dopamine
MOUSE MONOCLONAL

Catalog Number: AB-T11
Quantity: 50 microliters
Format: Lyophilized and reconstituted with deionized water / 50% glycerol
Host: Mouse
Isotype: IgG1, Kappa
Clone: HYDA001
Immunogen: Synthetic dopamine conjugated to bovine serum albumin

Specificity and Preparation:

Monoclonal antibody was obtained after BALB/c mouse immunization with the conjugates: Dopamine-Carrier proteins and hybridization of spleen cells with the myeloma cell line SP2/O/Ag14. Ascites production was performed in BALB/c mice. The ascites fluid was purified by ammonium sulfate precipitation.

This antibody targets conjugated dopamine. **This antibody does not recognize free dopamine.**

Using a conjugate Dopamine-Glutaraldehyde-BSA, antibody specificity was performed with an ELISA test by competition experiments with the following compounds:

COMPOUND	CROSS REACTIVITY §
Dopamine-G-BSA	1
L-DOPA-G-(Pc)	1/10,000
Tyrosine-G-(Pc)	1/36,000
Tyramine-G-(Pc)	1/>50,000
Noradrenaline-G-(Pc)	1/>50,000
Octopamine-G-(Pc)	1/>50,000
Adrenaline-G-(Pc)	1/>50,000
Dopamine	1/>50,000

G = Glutaraldehyde, BSA = bovine serum albumin

§ Dopamine-G-BSA concentration/unconjugated or conjugated catecholamine concentration at half displacement.

Usage and Storage:

Applications include ELISA (1/1,000-1/5,000); immunocytochemistry; immunohistochemistry (1/1,000-1/5,000); and immunoblotting (western blot, 1/1,000-1/2,000). Fixation of tissue for use with these antibodies should be done with glutaraldehyde. The use of paraformaldehyde in conjunction with glutaraldehyde may improve staining results. Store the antibody at 4°C for one month or -20°C in undiluted aliquots for up to one year. Avoid repeated freezing and thawing. Gently spin down material before use; 5-10 seconds in a microfuge should be adequate.

Available Control(s): Dopamine-G-BSA

References:

1. Goldman-Rakic PS, Leranath C, Williams SM, Mons N, Geffard M. (1989) Dopamine synaptic complex with pyramidal neurons in primate cerebral cortex. *Proc Natl Acad Sci USA* 86:9015-9019.
2. Schürmann FW, Elekes K, Geffard M. (1989) Dopamine-like immunoreactivity in the bee brain. *Cell Tissue Res* 256:399-410.

To view protocol(s) for this and other products please visit: www.ATSBio.com/support/protocols