

Swine IgM (mu) Rabbit Polyclonal
ATS-SELECT SECONDARY ANTIBODY

Catalog Number: AS-064
Quantity: 2 milliliters
Format: Antiserum, Lyophilized
Host: Rabbit
Immunogen: Swine IgM mu heavy chain

Background: Anti-Swine IgM antibody generated in rabbit specifically detects swine IgM. Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking 5 immunoglobulins together, the approximate molecular weight of IgM is 900kDa and possesses 10 binding sites (though due to the size of most antigens, not all sites are capable of binding at once). Due to this large size, IgM is typically isolated to the serum. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. Anti-Swine IgM antibody is ideal for investigators in Immunology, Microbiology, and Cell Biology.

Specificity & Preparation: This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against Swine IgM and Swine Serum. No reaction was observed against Swine IgG.

Usage: Anti-Swine IgM antibody are ideal for ELISA, western blotting, Immunohistochemistry, as well as other antibody detection methods.

ELISA 1:20,000 - 1:100,000

Immunohistochemistry 1:1,000 - 1:5,000

Western Blot 1:2,000 - 1:10,000

Working dilutions must be determined by end user.

Storage: Restore with deionized water (or equivalent), 2.0 mL. Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of receipt.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols