

**Armenian Hamster IgG (H&L) Secondary Goat Polyclonal
ATS-SELECT SECONDARY ANTIBODY**

Catalog Number: AS-069
Quantity: 2 milliliters
Format: Antiserum, Lyophilized
Host: Goat
Immunogen: Armenian Hamster IgG whole molecule

Background: Anti-Armenian Hamster IgG Antibody generated in goat detects Armenian Hamster IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the complement cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Specificity & Preparation: This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against Armenian Hamster IgG and Armenian Hamster Serum. Minimal cross reactivity is expected against golden syrian hamster IgG.

Usage: Secondary antibody to hamster IgG is available in a variety of formats. Anti-IgG Secondary Antibody is suitable for western blot, ELISA, ChIP and immunohistochemistry as well as other antibody based assays requiring lot-to-lot consistency.

ELISA 1:20,000 - 1:100,000

ChIP User Optimized

Immunohistochemistry 1:1,000 - 1:5,000

Western Blot 1:2,000 - 1:10,000

Working dilutions must be determined by end user.

Storage: Restore with deionized water (or equivalent), 2.0 mL. Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of receipt.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols