

**Human IgG F(ab')₂ Rabbit Polyclonal
ATS-SELECT SECONDARY ANTIBODY**

Catalog Number: AS-110
Quantity: 50 milligrams
Format: IgG, Lyophilized
Host: Rabbit
Immunogen: Human IgG F(ab')₂ fragment

Background: Anti-Human IgG F(ab')₂ Antibody generated in rabbit recognizes the dimeric Fab portion of the human IgG molecule. Human IgG F(ab')₂ is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled conditions of temperature, time and pH. F(ab')₂ molecules lack the Fc portion of IgG and therefore receptors that bind human IgG F(c) will not bind human IgG F(ab')₂ molecules. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Specificity & Preparation: This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Human IgG, Human IgG F(ab')₂ and Human Serum. No reaction was observed against Human IgG F(c).

Usage: Anti-Human IgG F(ab')₂ antibody is suitable for ELISA, western blot, and immunohistochemistry, as well as other assays requiring lot-to-lot consistency.

ELISA 1:20,000 - 1:100,000

Immunohistochemistry 1:1,000 - 1:5,000

Western Blot 1:2,000 - 1:10,000

Working dilutions must be determined by end user.

Storage: Restore with deionized water (or equivalent), 5.0 mL. Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of receipt.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols