

F(ab')₂ Mouse IgG (H&L) Goat Polyclonal
ATS-SELECT SECONDARY ANTIBODY

Catalog Number: AS-149
Quantity: 20 milligrams
Format: IgG F(ab')₂, Lyophilized
Host: Goat
Immunogen: Mouse IgG whole molecule

Background: F(ab')₂ Anti-Mouse IgG Antibody generated in goat was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab')₂ fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab')₂ fragments penetrate tissue samples and show better antigen recognition and signal generation in IHC. F(ab')₂ fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')₂ Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

Specificity & Preparation: This product is a F(ab')₂ fragment of IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation, ion exchange chromatography and pepsin digestion followed by chromatographic separation and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Mouse IgG and Mouse Serum. No reaction was observed against anti-Goat IgG F(c) or anti-Pepsin.

Usage: Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10⁶ cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.

ELISA 1:20,000 - 1:100,000

Immunohistochemistry 1:1,000 - 1:5,000

Western Blot 1:2,000 - 1:10,000

Working dilutions must be determined by end user.

Storage: Restore with deionized water (or equivalent), 2.0 mL. Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of receipt.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols