

Horse IgG (H&L) Goat Polyclonal
ATS-SELECT SECONDARY ANTIBODY

Catalog Number: AS-202
Quantity: 10 milligrams
Format: IgG, Liquid (sterile filtered)
Host: Goat
Immunogen: Horse IgG whole molecule

Background: Anti-Horse IgG Antibody generated in goat detects horse IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the complement cascade, and opsonization for phagocytosis. The whole IgG molecule possesses both the F(c) region, recognized by high-affinity Fc receptor proteins, as well as the F(ab) region possessing the epitope-recognition site. Both heavy and light chains of the antibody molecule are present. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Specificity & Preparation: This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Horse IgG and Horse Serum.

Usage: Anti-Horse IgG antibody has been tested by ELISA and is suitable for western blot and immunohistochemistry, as well as other assays requiring lot-to-lot consistency.

ELISA 1:20,000 - 1:100,000

Immunohistochemistry 1:1,000 - 1:5,000

Western Blot 1:2,000 - 1:10,000

Working dilutions must be determined by end user.

Storage: Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Expiration date is one (1) year from date of receipt.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols