Advanced Targeting Systems

1930 KELLOGG AVENUE
CARLSBAD, CA 92008 USA
01.858.642.1988 • WWW.ATSBIO.COM

Human IgG (H&L) Mouse Polyclonal
ATS-SELECT SECONDARY ANTIBODY

Catalog Number: AS-220
Quantity: 1 milligram
Format: IgG, Liquid (sterile filtered)
Host: Mouse
Immunogen: Human IgG whole molecule

Background: Anti-Human IgG (H&L) generated in mouse detects human Immunoglobulin G (IgG), both heavy and light chains of the antibody molecule are present. It is a protein complex composed of four peptide chains — two identical heavy chains and two identical light chains arranged in a Y-shape typical of antibody monomers. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobulins in humans, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Specificity & Preparation: This product was prepared from monospecific polyclonal ascites by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Mouse Serum, Human IgG and Human Serum.

Usage: Mouse Anti-Human IgG (H&L) has been tested by ELISA and western blot and is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user.
ELISA 1:5,000 - 1:50,000
Immunohistochemistry 1:1,000 - 1:5,000
Western Blot 1:500 - 1:5,000
Working dilutions must be determined by end user.

Storage: Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Expiration date is one (1) year from date of receipt.

To view protocol(s) for this and other products please visit: www.ATSbio.com/library/protocols