

Mouse IgG Fc Goat Polyclonal
ATS-SELECT SECONDARY ANTIBODY

Catalog Number: AS-243
Quantity: 2 milligrams
Format: IgG, Liquid (sterile filtered)
Host: Goat
Immunogen: Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat.

Background: Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of mouse IgG and often this fragment is removed from immunoglobulins to minimize receptor binding and lower background reactivity.

Specificity & Preparation: This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against Mouse IgG F(ab).

Usage: Anti-Mouse IgG F(c) fragment has been tested by ELISA, dot blot, and western blot and is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the end user.

ELISA 1:20,000 - 1:50,000

Immunohistochemistry 1:1,000 - 1:5,000

Western Blot 1:2,000 - 1:10,000

Working dilutions must be determined by end user.

Storage: Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Expiration date is one (1) year from date of receipt.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols