

**F(ab')₂ Mouse IgG F(ab')₂ Goat Polyclonal Pre-Adsorbed
ATS-SELECT SECONDARY ANTIBODY**

Catalog Number: AS-341
Quantity: 1 milligram
Format: IgG F(ab')₂, Liquid (sterile filtered)
Host: Goat
Immunogen: Mouse IgG F(ab')₂ fragment

Background: F(ab')₂ Anti-Mouse IgG F(ab')₂ Antibody generated in goat detects Mouse F(ab')₂. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(ab')₂ Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

Specificity & Preparation: This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Mouse IgG, Mouse IgG F(ab')₂ and Mouse Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), Mouse IgG F(c) or Bovine, Horse, Human, Rabbit, Rat and Sheep Serum Proteins.

Usage: F(ab')₂ Anti-Mouse IgG F(ab')₂ Antibody has been tested by ELISA and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10⁶ cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.
ELISA 1:20,000 - 1:100,000
Immunohistochemistry 1:1,000 - 1:5,000
Western Blot 1:2,000 - 1:10,000
Working dilutions must be determined by end user.

Storage: Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Expiration date is one (1) year from date of receipt.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols