

**Fab Rabbit IgG (H&L) Goat Polyclonal
ATS-SELECT SECONDARY ANTIBODY**

Catalog Number: AS-374
Quantity: 1 milligram
Format: IgG Fab, Liquid (sterile filtered)
Host: Goat
Immunogen: Rabbit IgG whole molecule

Background: Fab Anti-Rabbit IgG (H&L) Antibody generated in goat detects immunoglobulin g from rabbit, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.

Specificity & Preparation: This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum. No reaction was observed against anti-Papain or anti-Goat IgG F(c).

Usage: Fab Anti-Rabbit IgG (H&L) Antibody has been tested by dot blot and is suitable for highly specific immunological methods requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.
ELISA 1:4,000 - 1:20,000
Immunohistochemistry 1:1,000 - 1:5,000
Western Blot 1:2,000 - 1:10,000
Working dilutions must be determined by end user.

Storage: Store vial at 4° C prior to opening. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use. Expiration date is one (1) year from date of receipt.

To view protocol(s) for this and other products please visit: www.ATSBio.com/library/protocols