



6352 CORTE DEL ABETO, STE B  
CARLSBAD, CA 92011 USA  
01.858.642.1988 • WWW.ATSBIO.COM

**Alexa488-labeled Antibody to Angiotensin II receptor (AT-1BR), affinity-purified  
RABBIT POLYCLONAL**

**Catalog Number:** FL-N26AP  
**Quantity:** 50 micrograms  
**Format:** 50% PBS (0.14 M Sodium Chloride; 0.003 M Potassium Chloride; 0.002 M Potassium Phosphate; 0.01 M Sodium Phosphate; pH 7.4), 50% glycerol; no preservative.  
**Host:** Rabbit  
**Immunogen:** Synthetic peptide corresponding to a cytoplasmic region of the receptor conjugated to keyhole limpet hemocyanin (KLH)

**Background:**

The Angiotensin II type 1 receptor (AT-1R) is the primary effector of Angiotensin II, a key regulator of blood pressure and fluid homeostasis. It is involved in pathogenesis of several cardiovascular diseases such as hypertension, cardiac hypertrophy and congestive heart failure. Angiotensin II interacts with two types of G-protein coupled membrane receptors, AT-1R (type 1) and AT-2R (type 2). AT-1 has three isoforms in rat: AT-1A (359 aa), AT-1B (359 aa), and AT-1C (177 aa). Rat AT-1R's are predicted to contain seven transmembrane domains. The N-terminus is predicted to be extracellular, while the C-terminus is predicted to be cytoplasmic. AT-1R's are expressed in the liver, kidney, aorta, lung, uterus, ovary, spleen, heart, adrenal and vascular smooth muscle.

**Specificity and Preparation:**

This antibody recognizes the AT-1B isoform of the Angiotensin II type 1 receptor in rat, and does not recognize isoform AT-1A or AT-2. The antisera was generated in rabbits by immunization with a synthetic peptide corresponding to a cytoplasmic region of the receptor conjugated to keyhole limpet hemocyanin (KLH). Antisera was then affinity-purified by sequential passage through two affinity columns cross-linked with the carboxy terminal sequences of the rat AT-1B and AT-1A receptors. The antisera was retained by the first, but not the second column indicating recognition of the AT-1B receptor. It has been conjugated to the fluorescent dye Alexa488.

**Usage and Storage:**

Applications include immunolabeling (1:500).<sup>1,2</sup>

Gently spin down material before use; 5-10 seconds in a microfuge should be adequate. The material can be handled safely using normal laboratory precautions. See Lot Number for lot-specific storage instructions.

**References:**

1. Premer C, Lamondin C, Mitzey A, Speth RC, Brownfield MS. (2013) Immunohistochemical Localization of AT1a, AT1b, and AT2 Angiotensin II Receptor Subtypes in the Rat Adrenal, Pituitary, and Brain with a Perspective Commentary. *Int J Hypertens* 2013:175428.
2. Huang J, Hara Y, Anrather J, Speth RC, Iadecola C, Pickel VM (2003) Angiotensin II subtype 1A (AT1A) receptors in the rat sensory vagal complex: Subcellular localization and association with endogenous angiotensin. *Neuroscience* 122(1):21-36.
3. Speth RC, Grove KL, Brownfield MS (2001) Immunohistochemical localization of AT-1A and AT-1B angiotensin II receptor subtypes in the rat adrenal. *Endocrine Soc Mtg*, Denver CO, Abstract.

**To view protocol(s) for this and other products please visit: [www.ATSBio.com/support/protocols](http://www.ATSBio.com/support/protocols)**