



10451 ROSELLE STREET, #300, SAN DIEGO, CA 92121  
TELEPHONE (858) 642-1988 • FAX (858) 642-1989  
WWW.ATSBIO.COM • ATS@ATSBIO.COM

**Anti-Conjugated Tyrosine  
RABBIT POLYCLONAL  
AB-T030**

**Example of ELISA procol used to test conjugated tyrosine:**

1. Coating of conjugated tyrosine ( $15\mu\text{g/ml}$ ) in maxisorp well plates (Nunc) with a solution of sodium carbonate buffer  $0.05\text{M}$  (pH 9.6), during sixteen hours at  $4^{\circ}\text{C}$ .
2. Saturation of well plates with of a solution of PBS (pH 7.3) containing  $2.5\text{g/l}$  of BSA (Acros) and  $0.05\%$  Tween 20 (Acros) during one hour at  $37^{\circ}\text{C}$ .
3. Wash with PBS Tween (two times).
4. Anti-conjugated tyrosine antibodies will be diluted ( $1/1,000$ - $1/5,000$ ) in PBS containing  $2.5\text{g/l}$  BSA and  $10\%$  of glycerol,  $200\mu\text{l}$  by well plate (incubating during 2 hours at  $37^{\circ}\text{C}$ ).
5. Wash with PBS Tween (three times).
6.  $200\mu\text{l}$  of peroxidase-labeled sheep anti-rabbit (Bio-Rad) diluted ( $1/10,000$ ) in a solution of PBS containing  $2.5\text{g/l}$  BSA,  $10\%$  of glycerol and  $0.5\%$  of Tween will be applied by well plate (during one hour at  $37^{\circ}\text{C}$ ).
7. Well plates will be rinsed with a PBS Tween (three times).
8. And finally the peroxidase will be developed by incubating  $200\mu\text{l}$  by well plate of a citrate  $0.1\text{M}$ /phosphate  $0.2\text{M}$  (pH 5) solution containing  $0.4\%$  of OPD (Sigma) and  $0.03\%$  of hydrogen peroxide (Acros) for ten minutes in the dark, after that, we will stop the reaction by the addition of  $50\mu\text{l}$  of  $2\text{M}$  HCl.
9. The optical density will be measured at  $492\text{nm}$ , to obtain the different values.